


DUO-O

- Newly diagnosed advanced high grade epithelial OvC stage III-IV
 - Primary surgery or interval debulking surgery (IDS)
 - All comers
- N = 927



Stratification:

- 1) No residual vs. residual or IDS
- 2) tBRCAwt vs. tBRCAmut
- 3) Region: US; EU, RoW

Primary:

- PFS by Inv (BICR sensitivity)


Formal Secondary (tested):

- OS

Other secondary


- PFS2
- ORR
- HRQOL
- TFST, TSST, TDT

OReO design



Study design

ANITA




Stratification factors:

- Platinum based regimen selected
- PFI (6-12 months vs. > 12 months)
- BRCA mutation status (mutated vs. non-mutated)
- PD-L1 expression status (positive vs. negative vs. non informative)

Disegno dello studio

NEWTON



Fattori di minimizzazione per la parte randomizzata:

- sensibilità al platino (da 6 a <12 mesi vs ≥12 mesi);
- l'utilizzo di bevacizumab concomitante alla penultima linea di terapia a base di platino;
- la migliore risposta (parziale o completa) durante l'ultima linea di trattamento con platino.

IMN ISTITUTO DI RICERCHE
FARMACOLOGICHE
MARIO NEGRI - IRCCS

N.B. non arrotondare il valore del peso

Es. peso di 57.9 kg → gruppo non randomizzato